§ 180.484

§ 180.484 Flutolanil; tolerances for residues

(a) General. Tolerances are established for residues of flutolanil, N-(3-(1methylethoxy) phenyl)-2-(trifluoromethyl)benzamide, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only flutolanil and its metabolites converted to (trifluoromethyl) benzoic acid and calculated as flutolanil, in or on the following commodities:

Commodity	Parts per million
Cattle, fat	0.10
Cattle, kidney	1.00
Cattle, liver	2.00
Cattle, meat byproducts	0.05
Cattle, meat	0.05
Cotton, gin byproducts	0.20
Cotton, undelinted seed	0.20
Egg	0.05
Goat, fat	0.10
Goat, kidney	1.00
Goat, liver	2.00
Goat, meat byproducts	0.05
Goat, meat	0.05
Hog, fat	0.10
Hog, kidney	1.00
Hog, liver	2.00
Hog, meat byproducts	0.05
Hog, meat	0.05
Horse, fat	0.10
Horse, kidney	1.00
Horse, liver	2.00
Horse, meat byproducts	0.05
Horse, meat	0.05
Milk	0.05
Peanut	0.5
Peanut, hay	15.0
Peanut, meal	1.0
Potato	0.20
Potato, wet peel	0.20
Poultry, fat	0.05
Poultry, meat	0.05
Poultry, meat byproducts	0.05
Rice, bran	10.03
Rice, grain	7.0
Rice, hulls	25.0
Rice, straw	10.0
Sheep, fat	0.10
Sheep, kidney	1.00
Sheep, liver	2.00
Sheep, meat hyprodusts	0.05 0.05
Sheep, meat byproducts	
Soybean, forage	8.0
Soybean, hay	2.5
Soybean, seed	0.20 0.1
Turnip, greens	0.1
Vegetable, brassica, leafy, group 5	0.1

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]

(d) Indirect or inadvertent residues. Tolerances are established for the indirect or inadvertent residues of flutolanil, N-(3-(1-methylethoxy)phenyl)-2-

(trifluoromethyl)benzamide, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only flutolanil and its metabolites converted to 2-(trifluoromethyl) benzoic acid and calculated as flutolanil, in or on the following commodities.

Commodity	Parts per million
Wheat, bran Wheat, forage Wheat, grain Wheat hay Wheat, hay Wheat, straw	0.20 2.5 0.05 1.2 0.20

[60 FR 42458, Aug. 16, 1995, as amended at 61 FR 33044, June 26, 1996; 63 FR 42256, 42257, Aug. 7, 1998; 66 FR 10825, Feb. 20, 2001; 71 FR 7418, Dec. 13, 2006; 72 FR 35665, June 29, 2007; 73 FR 33017, June 11, 2008; 75 FR 17570, Apr. 7, 2010; 75 FR 80350, Dec. 22, 2010]

§ 180.485 Cyproconazole; tolerances for residues.

(a) General. (1) Tolerances are established for residues of the free and conjugated forms of the fungicide cyproconazole, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the proposed tolerance levels specified below is to be determined by measuring only cyproconazole (α -(4-chlorophenyl)- α -(1-cyclopropylethyl)-1H-1,2,4-triazole-1-ethanol) in or on the following commodities:

Commodity	Parts per million
Aspirated grain fractions	2.5
Cattle, fat	0.01
Cattle, meat byproducts (except liver)	0.01
Coffee bean, green (Imported) 1	0.1
Corn, field, forage	0.60
Corn, field, grain	0.01
Corn, field, stover	1.2
Goat, fat	0.01
Goat, meat byproducts (except liver)	0.01
Horse, fat	0.01
Horse, meat byproducts (except liver)	0.01
Peanut	0.01
Peanut, hay	6.0
Sheep, fat	0.01
Sheep, meat byproducts (except liver)	0.01
Soybean, forage	1.0
Soybean, hay	3.0

Environmental Protection Agency

Commodity	Parts per million
Soybean, oil	0.10 0.05 0.80 0.05 0.10 1.3 0.90

¹There are no U.S. registrations as of February 15, 2008 for use on coffee bean.

(2) A tolerance is established for the combined residues of the free and conjugated forms of the fungicide cyproconazole, including its metabolites and degradates, in or on the commodity in the table below. Compliance with the tolerance level specified below is to be determined by measuring only the sum of cyproconazole $(\alpha - (4$ chlorophenyl)-α-(1-cyclopropylethyl)-1H-1,2,4-triazole-1-ethanol) and its me- δ -(4-chlorophenyl)- β , δ tabolite dihydroxy-γ-methyl-1*H*-1,2,4-triazole-1hexenoic acid, calculated as the stoichiometric equivalent cyproconazole, in or on the following commodity:

Commodity	Parts per million
Milk	0.02

(3) Tolerances are established for the combined residues of the free and conjugated forms of the fungicide cyproconazole, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance level specified below is to be determined by measuring only the sum of cyproconazole (α -(4chlorophenyl)-α-(1-cyclopropylethyl)-1H-1,2,4-triazole-1-ethanol) and its me-2-(4-chlorophenyl)-3cyclopropyl-1-[1,2,4]triazol-1-yl-butane-2,3-diol, calculated as the stoichiometric equivalent of cyproconazole, in or on the following commodities:

Commodity	Parts per million
Cattle, liver Goat, liver Hog, liver Hog, liver Horse, liver Sheep, liver	0.50 0.50 0.01 0.50 0.50

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]

(d) Indirect or inadvertent residues. [Reserved]

[63 FR 53835, Oct. 7, 1998, as amended at 71 FR 71058, Dec. 8, 2006; 73 FR 27760, May 14, 2008; 78 FR 37473, June 21, 2013]

§ 180.486 Chlorethoxyfos; tolerances for residues.

(a) General. Tolerances are established for residues of the insecticide chlorethoxyfos, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only chlorethoxyfos, O,O-diethyl O-(1,2,2,2-tetrachloroethyl) phosphorothioate, in or on the commodity.

Commodity	Parts per million
Corn, field, forage	0.01
Corn, field, grain	0.01
Corn, field, stover	0.01
Corn, pop, grain	0.01
Corn, pop, stover	0.01
Corn, sweet, forage	0.01
Corn, sweet, kernel plus cob with husks re-	
moved	0.01
Corn, sweet, stover	0.01

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[76 FR 23498, Apr. 27, 2011]

§ 180.487 Pyrithiobac sodium; tolerances for residues.

(a) *General*. Tolerances are established for residues of the herbicide, pyrithiobac sodium, (sodium 2-chloro-6-[(4,6-dimethoxypyrimidin-2-

yl)thio]benzoate), resulting from the application of the pesticide chemical in or on the following foods/feeds:

Commodity	Parts per million
Cotton, gin byproducts	0.15 0.02

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]